

Received: July 2018

Accepted: August 2018

Epidemiologic Study of Patients with Cancer Admitted to Intensive Care

Roghayeh Honarjoo¹, Vida Shafipour², Mahmood Moosazadeh³, Ravanbakhsh Esmaeili^{4,*}

A B S T R A C T

9

1. Graduate Student of Intensive Care Nursing, Nasibe Sari School of Nursing and Midwifery, Student Research Committee of Mazandaran University of Medical Sciences, Sari, Iran.
2. Nasibeh Nursing and Midwifery Faculty, Mazandaran University of Medical Sciences, Sari, Iran.
3. Health Sciences Research center, Addiction Institute, Mazandaran University of Medical Sciences, Sari, Iran.
4. Assistant Professor, Faculty Member, Orthopedic Research Center, Mazandaran University of Medical Sciences, Sari, Iran.

*Corresponding Author:

Ravanbakhsh Esmaeili
Nasibeh Sari School of Nursing and Midwifery, Vesal Shirazi St, Sari
Tel: (+98)9111519435
Email: r.esmaeili90@gmail.com

Background: Cancer is the second leading cause of death in the world and the third cause of death in Iran. Evidence suggests that some cancer patients are admitted to intensive care units at their own request. The need for hospitalization in this ward can vary based on different factors such as gender, age, type of disease, etc. Therefore, this study has been designed in order to investigate the epidemiology of cancer patients admitted to intensive care units.

Methods: This is a descriptive cross-sectional study. The study population includes all cancer patients admitted to the intensive care units of Mazandaran University of Medical Sciences in the years 2013-2016. The records of all patients were included in the study through census and the necessary information was extracted. Data was analyzed using SPSS software version 24 in terms of percentage, mean, standard deviation through Chi-square test.

Results: 12273 patients were admitted to intensive care units during the study period, among which 380 (3.6%) were cancer patients. The age range of patients with cancer was between 16 and 96 years old, with a mean value of 60.11 years and a standard deviation of 17.49, and the majority of patients (52.4%) were mal. The frequency of patients in terms of cancer type were as follows; 40% gastrointestinal cancer, 20% brain cancer, 9.2% lung cancer, 6.6% kidney cancer, 6.6% breast cancer, 5.5% blood cancer, and 11.8% other cancers. The mortality rate of cancer patients admitted to ICUs was 36.1%. The frequency of patients with gastrointestinal, lung, blood, kidney and brain cancers admitted to the intensive care units in males was significantly higher than females ($p < 0.001$).

Conclusion: The results of this study indicate that, despite the need for intensive care in cancer patients, the lack of beds in these wards may lead to cancer patients being admitted less frequently than necessary. Therefore these wards may need to take practical measures in this respect, such as the establishment of intensive care units specifically for cancer patients.

Keywords: Cancer, Neoplasms, Hospital, epidemiology



2018; 10(4):9-15

www.bccrjournal.com

INTRODUCTION:

Cancer is one of the five leading causes of death in the world. Cancer (after heart disease) is the second leading cause of death in the United States, and the third leading cause of death in Iran¹. With an increase in population and in environmental risks, we are faced with an increase in the incidence of cancer, despite the general improvement in health conditions, and in some societies malignant diseases have become the nation's top health priority. Increase in the incidence of cancer may be due to changes in lifestyle, the aging of the population and increased environmental risks². It is likely that this disease will be the major health issue of the next decade³ therefore, accurate registration of prevention and treatment programs and timely treatment are required⁴. In Iran, cancer claims over 30000 lives annually, and the incidence rate is expected to double in the next two decades⁵. According to the International Agency for Research on Cancer, the mortality rate caused by cancer in 2007 was estimated to be over 1.8 million people, and is estimated to reach 10 million deaths in 2030, with rates being highest in developing countries⁶. In 2012, about 14 million and in 2015 about 8.5 million cancer-related deaths were reported worldwide⁴.

The intensive care unit is a well-equipped medical and nursing department and has been recognized as one of the most sensitive parts of the hospital and the best place for delivering patient care⁷. Patients with different diagnoses are admitted in this ward⁸ and these wards are well suited for heart and respiratory monitoring of patients⁹. Admission to ICUs affects the success and failure of treatment¹⁰, and studies have shown that observing standards and admission to the intensive care unit increases the probability of survival¹¹.

It is noteworthy that intensive care units in many developing countries face many challenges¹². The number

of beds in these wards are limited¹³, posing a challenge for health care providers worldwide. In the study conducted by Aksoy et al. in Turkey¹⁴ and Monique et al. in the Netherlands over the past four years, 21.33% and 13.5% of patients admitted to ICUs were cancer patients, respectively¹⁵.

Considering the mentioned issues that indicate the role of intensive care in increasing the survival rate of cancer patients and improving the quality of care provided to these patients, in the future, it is expected to see an increase in demand and acceptance of these patients. This approach will create serious challenges due to the lack of resources and facilities needed to equip the intensive care units. For this purpose, it is necessary to plan and adopt an appropriate strategy for supplying and equipping intensive care units. In this regard, studying the status quo and the benefit gained among cancer patients from these wards is a priority. Our search of available sources and available databases, yielded no evidence regarding the status of hospitalization of patients with cancer in ICUs in Iran, and especially in northern Iran. The present study was designed and conducted to evaluate the epidemiology of cancer patients admitted to intensive care units, which can be used in future planning.

METHODS:

This study is a descriptive cross-sectional study. The study population consisted of all patients with cancer admitted to the intensive care units of educational hospitals affiliated to Mazandaran University of Medical Sciences between the years 2013-2016. Inclusion criteria were age greater than 16 years, cancer diagnosis based on pathological results and ICU admission. The source of data collection were patient hospital files. Sampling method was census, since the records of all patients with cancer admitted to the intensive care units of the educational hospitals affiliated to Mazandaran

University of Medical Sciences were investigated during the research period.

After approval of the project by the Research Council and the Ethics Committee in the Medical Biology Research Department of the university and after obtaining a license from the Deputy of Research and Technology of Mazandaran University of Medical Sciences, permission was obtained from the hospital authorities, and one of the members of the research team visited the medical records sectors of selected hospitals (Imam Sari, Bouali Sari and Razi Ghaemshahr) in order to collect the data. A checklist was developed by the research team using documentation, scientific literature and the opinions of medical and nursing experts in order to record and collect data. Afterwards, content validity was assessed and verified by 10 professors of nursing and intensive care. Data analysis was done in SPSS version 24. After refining the data and removing possible defects, according to the type of study, frequency percentage, mean, standard deviation, median, minimum and maximum were used to describe the variables. Also, Chi-square test was used for comparison. The criterion for judgment

was a significance level less than 0.05.

RESULTS:

The results of this study showed that of the 12273 patients hospitalized in intensive care units during the years 2013 to 2016, 380 (3.09%) of the patients were diagnosed with cancer. Of these 380 patients, 47.6% were female and 52.4% were male. The age range of patients was 16 to 96 years old with an average age of 60.11 years and a standard deviation of 17.49. The frequency of patients aged less than 70 years was 68.2% and older than 70 years was 31.8%.

Table 1 shows that the frequency of patients with gastrointestinal, brain, lung, kidney, breast, blood and other cancers were respectively 40%, 20%, 9.2%, 6.6%, 6.6%, 5.5% and 11.8%. The gender ratio of males to females in patients with gastrointestinal, brain, lung, kidney, breast, blood and other cancers admitted to intensive care units was 1.26, 1.16, 1.69, 1.33, 2.24, 0.04 and 0.79 respectively ($P < 0.001$). Also, as is shown in **Table 2**, type of cancer was significantly related to metastasis, place of living, result of admission, history of smoking, history of admission, reason for admission

Table 1. Comparison of the frequency of cancer type in terms of gender in the intensive care unit

Type of cancer	Total number (%)	Gender –n (%)		Ratio male/female	P
		Male	Female		
<i>Digestive</i>	152 (40)	85 (55.9)	67 (44.1)	1.26	<0.001
<i>Brain</i>	20 (76)	41 (53.9)	35 (19.3)	1.16	
<i>Lung</i>	35 (9.2)	22 (62.9)	13 (37.1)	1.69	
<i>Blood</i>	21 (5.5)	12 (57.1)	9 (42.9)	1.33	
<i>Renal</i>	26 (6.8)	18 (69.2)	8 (30.8)	2.24	
<i>Breast</i>	25 (6.6)	1 (4)	24 (96)	0.041	
<i>Other</i>	45 (11.8)	20 (44.4)	25 (55.6)	0.79	
<i>Total</i>	380 (100)	199 (52.4)	181 (47.6)	1.09	

Table 2. Comparison of the status of patients with cancer at the time of admission to the intensive care unit based on different variables

Variable	Digestive number (%)	Lung number (%)	Blood number (%)	Breast number (%)	Renal number (%)	Brain number (%)	Other number (%)	P
Metastasis	Yes	37 24.5	10 29.4	2 9.5	18 72	11 42.3	14 31.1	0.00
	No	114 75.5	24 70.6	19 90.5	7 28	15 57.7	31 85.1	
Reason for admission	Monitoring	142 93.4	28 80	17 81	24 96	24 92.3	36 93.4	80
	Complications of metastasis	2 1.3	0 0	0 0	1 4	0 0	0 0	0 0
	Complication medicine	0 0	0 0	0 0	0 0	0 0	1 1.3	0 0
	Resuscitation	8 5.3	7 20	4 19	0 0	2 7.7	4 5.3	9 20
History of smoking	Yes	46 31.7	18 51.4	5 23.8	2 8	10 38.5	14 18.9	11 25.6
	No	99 68.3	17 48.6	16 76.2	23 92	16 61.5	60 81.1	32 74.4
Feeding method at time of admission	Oral	2 1.3	5 14.3	2 9.5	3 12	2 7.7	3 3.9	3 6.7
	Gavagh	2 1.3	0 0	0 0	3 12	0 0	0 0	0 0
	Venous	147 96.7	29 82.9	18 85.7	19 76	24 92.3	72 94.7	41 91.1
	Oral venous	1 0.7	1 2.9	0 0	0 0	0 0	1 1.3	0 0
	Vevousgavagh	0 0	0 0	1 4.8	0 0	0 0	0 0	1 2.2
	Death	53 34.9	20 57.1	12 57.1	13 52	8 30.8	16 21.1	15 33.3
Result of admission	Move to unite	96 63.2	14 40	7 33.3	12 48	17 65.4	29 78.9	29 64.4
	Personal satisfaction	3 2	1 2.9	0 0	0 0	1 3.8	0 0	1 2.2
History of admission	Another hospital	0 0	0 0	2 9.5	0 0	0 0	0 0	0 0
	Yes	50 33.6	21 65.6	5 26.3	8 32	11 45.8	26 34.2	13 29.5
Place of living	No	99 66.4	11 34.4	14 73.7	17 68	13 54.2	50 65.8	31 70.5
	City	80 52.6	18 51.4	14 66.7	18 72	12 46.2	28 36.8	26 57.8
Roly	72 47.4	17 48.6	7 33.3	7 28	14 53.8	48 63.2	19 42.2	

and feeding method at the time of admission of the patient to the intensive care unit ($P < 0.05$).

Also, the results of this study showed that among patients with cancer who were hospitalized in intensive care units, 35.3% had a previous history of admission to these wards and 66.1% of them received drugs during admission. Results also showed that 9.2% had a familial history of cancer and 1.8% had a history of CPR before hospitalization.

The results of this study indicate that 54.7% of patients had a previous history of surgery and 34.5% of them had been afflicted with bedsores while hospitalized in intensive care units. Patients with gastrointestinal cancer had the highest percentage of bedsores during hospitalization in intensive care units. Findings also showed that 49.2% of patients used mechanical ventilation during hospitalization and 90% of these patients were admitted to these wards due to their need for appropriate monitoring of vital signs. At the time of entry of cancer patients into the intensive care unit, intravenous feeding among patients with gastrointestinal, kidney, lung, blood, breast, brain cancers was respectively, 96.7%, 92.3%, 82.9%, 85%, 76%, 49.7%. The outcome of hospitalization of these patients in the intensive care units included 61.6% being transferred to wards, 36.1% mortality, 1.6% being discharged with personal consent and 0.5% being transferred to other hospitals. Also, the findings showed that in terms of level of consciousness and Glasgow Coma Scale, most patients had a GCS in the range of ¹³⁻¹⁵.

The results of this study also indicated that surgery (64.4%) and palliative care (27.4%) were the most commonly used treatments in these patients, and the most common underlying diseases in these patients were pulmonary disease, anemia, and diabetes, in that order. The results of the study also indicated that factors affecting the mortality of patients with cancer admitted in intensive care units using single-variable and multiple variables of logistic regression include, mechanical

ventilation, type of cancer, type of treatment, family history, smoking, underlying diseases and age of the patient, and that probability of death was lower among patients receiving mechanical ventilation.

DISCUSSION:

The results of this study showed that 3.9% of patients admitted to intensive care units were patients with various types of cancers. Whereas in the studies conducted by Aksoy et al. (2016) in Turkey, lasting two years¹⁴, Monique et al. (2012) in the Netherlands lasting four years¹⁵, Jonge and Bos (2009) in Europe¹⁶, Xia and Wang (2016) in China over a period of 3 years¹⁷, and in the study of Hawari et al. (2016) in Jordan over five years¹⁸, respectively 21.03%, 13.5%, 15%, 17% and 22.6% of patients admitted to intensive care units had cancer. The results of the aforementioned studies are different from the results gathered in this study. Concerning this difference, it can be said that in Iran, unfortunately, due to the lack of intensive care beds, less cancer patients are admitted to the intensive care unit, while in other countries it is possible to take care of these patients in intensive care units and more patients will benefit from intensive care services. According to the findings of this study, the most common cancer among both genders was gastrointestinal cancer with a frequency of 44.1% in females and 55.9% in males. These findings are consistent with the results obtained by Aksoy et al.¹⁴ and Xia and Wang¹⁷ And also with the results of the registry in the two northern provinces of our country^{19,20}.

The study also found that admitted cancer patients were more commonly male, which is similar to the results obtained by Aksoy et al.¹⁴ and Xia and Wang¹⁷. Therefore, the results of other studies and the current study confirm that, general male cancer patients are more frequently admitted to ICUs than female. In this study, as in the study by Aksoy et al.¹⁴, Bird et al.²¹ and Xia A and Wang¹⁷, it was indicated that, the

majority of patients with cancer hospitalized in intensive care units were approximately 70 years old. The reason for this can be that, with new advances in the treatment and diagnosis of cancer, this disease is diagnosed at an earlier age¹⁴.

Also, according to the results of this study 61.8% of cancer patients were transferred to public wards following treatment in intensive care units. In the study by Aksoy et al.¹⁴, and Zafra et al.²², respectively 53.3% and 68 % of these patients were transferred to the general ward. It seems that this difference may be associated with hospital rules and regulations in different countries. The findings of the recent study also show that, surgery was the most commonly used treatment method in patients in the intensive care units, a finding which is consistent with the results obtained by Aksoy et al.¹⁴ and Xia and Wang¹⁷ and Soares et al.²³. Given that today most cancers are diagnosed in a timely manner and surgery is used for treatment, therefore, surgery is the most commonly used treatment in all stages of cancer and it is done in a variety of different ways and depends on the discretion of doctors²⁴.

While assessing the mortality rate of cancer patients admitted to the intensive care unit, the results of this study showed that 36.1% died, and that probability of death was lower among patients with mechanical ventilation. These results are consistent with the study conducted by Monique et al.¹⁵ and Hawari et al.¹⁸. But in the findings of Xia and Wang¹⁷, the mortality rate of cancer patients in the intensive care units was 14.9%. It seems that the use of new therapeutic interventions, including medical treatments, mechanical ventilation and alternative renal treatments, and also timely admission of these patients to intensive care units can reduce the mortality rate of these patients.

According to the results of our study, the highest mortality rate in cancer patients in these wards was related

to blood and lung cancer, respectively, which is consistent with the results obtained by Hawari et al. at a cancer center in Jordan over a period of 5 years¹⁸.

In this study, 49.2% of cancer patients admitted to intensive care units required mechanical ventilation which is similar to the results of Bird et al. (2012) who found that 51.9% of patients were subjected to mechanical ventilation²¹. This type of treatment is used to support the respiratory system²⁵. In recent decades, significant survival has been reported in cancer patients undergoing mechanical ventilation in intensive care units²⁶.

CONCLUSION:

Considering that in this study, the admission rate of cancer patients in ICUs was lower than in other countries, and that the intensive care provided in these wards increases the survival rate of these patients, and also considering the fact that there is a severe shortage of intensive care beds in Iran, (which is probably due to the fact that the admission percentage of cancer patients in these ward is lower, while these patients need intensive care for treatment and recovery), therefore, it is necessary to take certain measures for admission of these patients to intensive care units and allocate a suitable percentage of intensive care beds for these patients. This can be achieved by establishing intensive care units specifically for cancer patients. One of the strengths of this study is the review of a 4-year period of hospitalization of patients with cancer admitted to intensive care units, which has not previously been done in Iran. The use of a retrospective approach is one of the research constraints which prohibits accurate data assessment and follow-up. Therefore, we suggest that in the future long-term studies or prospective studies be carried out in relation to this research.

REFERENCES:

1. Aminzadeh A, Motaghi A, Mohammadi E. Epidemiologic

- study of oral and paraoral malignancies in one cancer referral center in Isfahan during a 5-year period. *Journal of Isfahan Dental School*. 2013;8(6).
2. Warnakulasuriya S. Global epidemiology of oral and oropharyngeal cancer. *Oral oncology*. 2009;45(4-5):309-16.
 3. Torre LA, Siegel RL, Ward EM, Jemal A. Global cancer incidence and mortality rates and trends—an update. *Cancer Epidemiology and Prevention Biomarkers*. 2015.
 4. van Leeuwen RW, Jansman FG, van den Bemt PM, de Man F, Piran F, Vincenten I, et al. Drug–drug interactions in patients treated for cancer: a prospective study on clinical interventions. *Annals of Oncology*. 2015;26(5):992-7.
 5. Jafari A, Najafi S, Moradi F, Kharazifard M, Khami M. Delay in the diagnosis and treatment of oral cancer. *Journal of Dentistry*. 2013;14(3):146.
 6. Jemal A, Center MM, DeSantis C, Ward EM. Global patterns of cancer incidence and mortality rates and trends. *Cancer Epidemiology and Prevention Biomarkers*. 2010;1055-9965. EPI-10-0437.
 7. Yousefinezhadi T, Mosadeghrad AM, Mohammad A, Ramezani M, SARI AA. An analysis of hospital accreditation policy in Iran. *Iranian Journal of Public Health*. 2017;46(10):1347.
 8. Fernández-Ortega J, Herrero Meseguer J, Martínez García P. Guidelines for specialized nutritional and metabolic support in the critically-ill patient. Update. Consensus SEMICYUC-SENPE: Indications, timing and routes of nutrient delivery. *Nutricion hospitalaria*. 2011;26(2).
 9. Günther A. Skin conductance variability and stressful exposures in critical care: Inst för fysiologi och farmakologi/Dept of Physiology and Pharmacology; 2016.
 10. Jarahzadeh MH, Emamimeybodi AH, Halvani A, Mirjalili MR, Farnia F. The Prognosis of Patients with Admission Hypomagnesemia in Intensive Care Unit Related to APACHE II, III and SOFA Scores. *scientific journal of ilam university of medical sciences*. 2017;25(1):45-53.
 11. Van Sambeek J, Cornelissen F, Bakker P, Krabbendam JJ. Models as instruments for optimizing hospital processes: a systematic review. *International journal of health care quality assurance*. 2010;23(4):356-77.
 12. Kwizera A, Dünser M, Nakibuuka J. National intensive care unit bed capacity and ICU patient characteristics in a low income country. *BMC research notes*. 2012;5(1):475.
 13. Aksoy Y, Kaydu A, Sahin OF, Kacar CK. Analysis of cancer patients admitted to intensive care unit. *Northern clinics of Istanbul*. 2016;3(3):217.
 14. Bos MM, de Keizer NF, Meynaar IA, Bakhshi-Raiez F, de Jonge E. Outcomes of cancer patients after unplanned admission to general intensive care units. *Acta oncologica*. 2012;51(7):897-905.
 15. de Jonge E, Bos MM. Patients with cancer on the ICU: the times they are changing. *Critical Care*. 2009;13(2):122.
 16. Xia R, Wang D. Intensive care unit prognostic factors in critically ill patients with advanced solid tumors: a 3-year retrospective study. *BMC cancer*. 2016;16(1):188.
 17. Hawari FI, Nazer LH, Addassi A, Rimawi D, Jamal K. Predictors of ICU admission in patients with cancer and the related characteristics and outcomes: a 5-year registry-based study. *Critical care medicine*. 2016;44(3):548-53.
 18. Shokrzadeh M, Mohammadpour A, Tabari YS, Parvizi Almani S. Investigating the Distribution of ERCC2 (rs13181) Gene Polymorphism in Gastric Cancer Patients in Mazandaran: A Case-control Study. *Journal of Genetic Resources*. 2017;3(1):54-60.
 19. Farvid MS, Malekshah AF, Pourshams A, Poustchi H, Sepanlou SG, Sharafkhan M, et al. Dairy food intake and all-cause, cardiovascular disease, and cancer mortality: The Golestan Cohort Study. *American journal of epidemiology*. 2017;185(8):697-711.
 20. Bird G, Farquhar-Smith P, Wigmore T, Potter M, Gruber P. Outcomes and prognostic factors in patients with haematological malignancy admitted to a specialist cancer intensive care unit: a 5 yr study. *British journal of anaesthesia*. 2012;108(3):452-9.
 21. Zafra M, Carrillo A, Vicente MA, Sánchez Cánovas M, Ivars Rubio A, Ballester Navarro I, et al. Long-term survival and performance status of cancer patients after discharge from ICU (intensive care unit). *American Society of Clinical Oncology*; 2018.
 22. Soares M, Caruso P, Silva E, Teles JM, Lobo SM, Friedman G, et al. Characteristics and outcomes of patients with cancer requiring admission to intensive care units: a prospective multicenter study. *Critical care medicine*. 2010;38(1):9-15.
 23. Schiavone MB, Moukarzel L, Leong K, Zhou QC, Afonso AM, Iasonos A, et al. Surgical site infection reduction bundle in patients with gynecologic cancer undergoing colon surgery. *Gynecologic oncology*. 2017;147(1):115-9.
 24. Girard TD, Alhazzani W, Kress JP, Ouellette DR, Schmidt GA, Truitt JD, et al. An official American Thoracic Society/American College of Chest Physicians Clinical Practice Guideline: liberation from mechanical ventilation in critically ill adults. Rehabilitation protocols, ventilator liberation protocols, and cuff leak tests. *American journal of respiratory and critical care medicine*. 2017;195(1):120-33.
 25. Saillard C, Mokart D, Lemiale V, Azoulay E. Mechanical ventilation in cancer patients. *Minerva Anesthesiol*. 2014;80(6):712-25.
 26. Saillard C, Mokart D, Lemiale V, Azoulay E. Mechanical ventilation in cancer patients. *Minerva Anesthesiol*. 2014;80(6):712-25.